

**Problem 1.** Consider the binary operation  $\sqcup$  defined on the set  $\mathbb{Z}$  of integers by the formula

$$m \sqcup n = |m - n|$$

**Part (a).** Is this operation commutative?

**Part (b).** Is this operation associative?

**Problem 2.** Consider the binary operation  $\cup$  defined on the set  $\mathbb{R}$  of real numbers by the formula

$$a \cup b = \sqrt[4]{a^4 + b^4}$$

**Part (a).** Is this operation commutative?

**Part (b).** Is this operation associative?

**Problem 3.** Consider the set  $\mathbb{Z}_{10} = \{0,1,2,3,4,5,6,7,8,9\}$  along with the function  $f : \mathbb{Z}_{10} \rightarrow \mathbb{Z}_{10}$  defined by  $f(x) = 4 \boxtimes_{10} x$ .

**Part (a).** Write the function  $f$  in tabular notation.

$$f : \begin{pmatrix} 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \end{pmatrix}$$

**Part (b).** Is this function a member of the family  $\mathcal{P}_{\mathbb{Z}_{10}}$ ?

**Problem 4.** Consider the following members of the permutation group  $\mathcal{P}_7$ .

$$f : \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 4 & 3 & 2 & 5 & 1 & 7 & 6 \end{pmatrix} \quad g : \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 3 & 4 & 5 & 2 & 6 & 1 & 7 \end{pmatrix}$$

**Part (a).** Construct the formulas for the permutations  $g^{-1}$ ,  $f \circ g$ , and  $g \circ f \circ g^{-1}$ .

**Part (b).** What is the smallest positive integer  $n$  such that  $f^n$  is the identity for  $\mathcal{P}_7$ ?

**Problem 5.** Consider the group  $\mathcal{Z}_{20} = (\mathbb{Z}_{20}, \boxplus_{20})$ .

**Part (a).** What is the smallest positive integer  $n$  such that  $8^n$  is the identity for  $\mathcal{Z}_{20}$ ?

**Part (b).** What is the largest negative integer  $m$  such that  $3^m$  is the identity for  $\mathcal{Z}_{20}$ ?

**Problem 6.** A binary operation on the set  $X = \{a, b, c, d, e\}$  has been partially defined by the table below. If we know that  $(X, *)$  is a Sudoku algebra, complete the table.

*	a	b	c	d	e
a	a	c	b	e	
b	b	a	e	d	c
c		d	a		e
d	e	b	d	c	a
e	d			a	b

**Problem 7.** Consider the set  $M = \{I, A, B, C, D, E\}$ , where

$$I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \quad A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad B = \begin{bmatrix} 0 & 1 \\ -1 & -1 \end{bmatrix}$$

$$C = \begin{bmatrix} -1 & -1 \\ 0 & 1 \end{bmatrix} \quad D = \begin{bmatrix} -1 & -1 \\ 1 & 0 \end{bmatrix} \quad E = \begin{bmatrix} 1 & 0 \\ -1 & -1 \end{bmatrix}$$

Recall from linear algebra that the product of two  $2 \times 2$  matrices is given by the formula

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} * \begin{bmatrix} w & x \\ y & z \end{bmatrix} = \begin{bmatrix} aw + by & ax + bz \\ cw + dy & cx + dz \end{bmatrix}$$

**Part (a).** Fill in the table below.

*	I	A	B	C	D	E
I						
A						
B						
C						
D						
E						

**Part (b).** Is  $(M, *)$  a Sudoku algebra? Explain.

**Part (c).** Is  $(M, *)$  a group? Explain.

**Problem 8.** Consider the set  $I = \{x \in \mathbb{R} : 0 < x \leq 1\}$  along with the relation

$$a \div b = \begin{cases} ab & \text{if } a < b \\ b/a & \text{if } b \leq a \end{cases}$$

**Part (a).** Show that this relation is a binary operation on the set  $I$ .

**Part (b).** Show that  $x = 2/3$  and  $x = 1/6$  are *both* solutions to the equation  $(1/2) \div x = 1/3$ .

**Part (c).** Show that  $\varepsilon = 1$  serves as an identity for the algebra  $(I, \div)$ .

**Part (d).** Show that every element of  $I$  serves as its own inverse with respect to the operation  $\div$ .

**Part (e).** Without using a counterexample, explain why the operation  $\div$  *cannot* be associative. Can you find a specific counterexample?

**Problem 9.** Suppose that  $\mathcal{X} = (X, *)$  is a group, and suppose we know that there exist  $a, b, c \in X$  such that

- $a * c = c * a$ ,
- $b * c = c * b$ , and
- $c = (a * b) * (b * a)^{-1}$ .

Use this information to show that  $b * (c * a) = a * b$ .

Suppose that  $\mathcal{X} = (X, *)$  is a group with identity  $\varepsilon$ , and suppose that  $a \in X$ . We say that  $a$  has *finite order* in this group provided there exists a smallest positive integer  $p$  such that  $a^p = \varepsilon$ .

**Problem 10.** Fill in the gaps in the proof of the following result.

*If  $\mathcal{X} = (X, *)$  is a finite group, then every member of  $X$  has finite order in the group  $\mathcal{X}$ .*

**Proof.** Let  $X = \{\varepsilon, a_1, \dots, a_n\}$ , where  $\varepsilon$  is the identity for  $\mathcal{X}$ . Let  $b \in X$  and consider the set

$$S_b = \{b^j : j \in \mathbb{Z}^+\}$$

There must exist positive integers  $j < k$  such that  $b^j = b^k$ . [WHY?]

Consequently, we know that there exists a positive integer  $m$  such that  $b^m = \varepsilon$ . [WHY?]

It follows that there is a smallest positive integer  $p$  such that  $b^p = \varepsilon$ . [WHY?]

We must therefore conclude that  $b$  has finite order in the group  $\mathcal{X}$ .

**Problem 11.** Suppose that  $\mathcal{X} = (X, *)$  is a group with identity  $\varepsilon$ , and suppose that  $a \in X$  has finite order  $p$ . Prove that the elements  $a^1, a^2, \dots, a^{p-1}, a^p$  are all distinct. (Hint: Suppose, for purposes of contradiction, that two of these elements are equal.)